

**IDENTIFICATION AND CHARACTERIZATION OF
RHIZOSPHERIC MICROBIAL ISOLATES WITH
ANTIMICROBIAL (ANTIFUNGAL) ACTIVITY
FROM *SORGASTRUM NUTANS***

MRRIDULA DANGI NARWAL

M.SC. (GENETICS)

M.TECH. (BIOTECHNOLOGY)

U.I.E.T.

MAHARSHI DAYANAND UNIVERSITY

ROHTAK

HARYANA

INDIA

Abstract

The goal of this review was to disconnect and describe a rhizospheric bacterium from *Sorgastrum nutans*, developing around the betray district of Rajasthan (India). Plant growth promoting rhizobacterias (PGPR) are known to impact plant development by different immediate or circuitous systems. Separated strain was tried for different PGP

characteristics like 1-aminocyclopropane-1-carboxylate (ACC) deaminase movement, phosphate solubilization, indole acidic corrosive generation, creation of siderophore, nitrogen obsession and alkali generation. Bio-control capacity of detach was screened by antagonistic action against certain parasitic/bacterial pathogens and additionally Hydrogen cyanide(HCN)

creation. Selected test strain was likewise biochemically characterized. Advance recognizable proof of identification was performed by PCR based 16S rRNA qualitative sequencing. Besides assessment of the segregate SNP-18 showing different plant growth promoting (PGP) qualities on soil/plant framework is on-going to reveal their viability as successful PGPR.

Key words: QPGPR, Rhizosphere, ACC deaminase, PCR, Sorgastrum nutans, Desert region of Rajasthan, India.

Introduction

Plant growth promoting rhizobacteria (PGPR) is a gathering of heterogeneous microbes that can be found in the rhizosphere, at root surfaces and in relationship with roots, which can enhance the degree or nature of plant development straightforwardly as well as by implication. Term PGPR was initially presented in 1978 by Kloepper and partners. The immediate advancement by PGPR happens regarding

giving the plant a plant development, advancing substances that is orchestrated by the bacterium or encouraging the take-up of certain plant supplements from nature. The backhanded advancement of plant development happens when PGPR keep the malicious impact of phytopathogenic microorganisms. In the most recent couple of decades, a huge gathering of microscopic organisms including types of Alcaligenes, Arthobacter, Azospirillum, Bacillus, Burkholderia, Pseudomonas, Klebsiella, Enterobacter, and Serratia has answered to upgrade plant development (Kloepper et al., 1989; Glick, 1995).

However the correct systems of PGPR interceded plant development are not completely seen, but rather are thought to happens through (i) the capacity to create plant development controllers, for example, indoleacetic corrosive, gibberellic corrosive, cytokinins (Glick, 1995), (ii) asymbiotic N₂ obsession (Boddey and Dobreiner, 1995), (iii) solubilization of mineral phosphates and different supplements (Gaur, 1990) (iv)

hostile movement against phytopathogenic microorganisms by generation of siderophores, anti-toxins and HCN generation (Scher and Baker, 1982). A large portion of the microscopic organisms of family *Pseudomonas* and *Bacillus* abused as biocontrol specialist. Some PGPR may advance plant development by implication by influencing advantageous N₂ fixation, nodulation or knob inhabitation (Fuhrmann and Wollum, 1989). Furthermore, plant development advancing bacterial strains must be rhizospheric skillful, for their survival and colonization in the rhizospheric soil (Cattelan et al., 1999). The current rural part are experiencing different ecological difficulties including atmosphere/climate conditions, abiotic stress conditions, attributes of soil or injurious action of the indigenous microbial greenery of the dirt (Bent et al., 2001). As per FAO (Food and Agriculture Organization of United States) 20-40% worldwide harvest creation extremely influenced each year because of the movement of different phytopathogenic microorganism. In addition, utilization of

pesticides antagonistically influencing the biological condition by expanding different sorts of contamination and extraordinary impact on soil fruitfulness. Consequently utilization of microbial inoculants as bio-control operators could forestalls plant pathogens through a polyphonic instrument like generation of anti-infective agents or lyric catalysts rivalry with other soil smaller scale living beings for supplements and initiated systemic resistance. Henceforth utilization of advantageous rhizospheric microbes could be utilized as a compelling contrasting option to substance manures, pesticides and other agrarian supplements (Ashrafuzzaman et al., 2009).

Also, microorganism with ACC deaminase action has an extra advantage of plants through decreasing the anxiety inducible ethylene level in host plants (Karthikeyan et al., 2012; Jha et al., 2012). The adequacy of ACC deaminase catalyst had been accounted for in different ecological burdens like flooding, warmth push, dry season, metal tainting, natural contaminations, injuring,

pathogen and creepy crawly disease (Glick et al., 2007; Morgan and Drew 1997). *Sorgastrum nutans*, generally known as yellow Indian grass, has a place with family Poaceae. Yellow Indian grass is local to prairie natural surroundings and tolerant to shade. In for sake of Rajasthan, it can be seen much of the time developing in rural grounds, consequently our point was to detach and describe the rhizospheric microorganisms related with *Sorgastrum nutans*.

Biochemical portrayal

Separate was screened for different biochemical tests, for example, Gram recoloring, starch agar test, IMViC. Exercises for a few compounds, for example, catalase, oxidase, lipase, amylase, protease, nitrate reductase was tried according to standard conventions (Prescott and Harley, 2002). The capacity of the separator to use different carbon sources was tried utilizing starch use test pack (KB 009, Himedia, India). Affectability of detach to

different anti-infective agents to be specific gentamicin, ampicillin, erythromycin, kanamycin, antibiotic medication, streptomycin, and chloramphenicol was tried by anti-microbial plates (HTM 002, Himedia India).

Sub-atomic ID and phylogenetic examination Isolation of genomic DNA

For disconnection of genomic DNA (g-DNA), an independent settlement was picked from the naturally developed culture of bacterial separate and was developed in Luria-soup medium (Himedia, India) for overnight at 37°C with shaking at 150 RPM. Add up to genomic DNA of strain was confined by Quiagen DNA segregation unit (Quiagen, USA).

PCR enhancement of 16S rRNA quality

ID of confine SNP-18 was finished by 16S rRNA quality sequencing. Genomic DNA of bacterial segregate was opened up with general preliminary 27F1 (52 - AGAGTTTGATCMTGGCTCAG-32) and

1494Rc (52 -
 TACGGCTACCTTGTTACGAC-32) in a
 25 µl response blend containing 10 X
 cushion (with 2.5mm MgCl₂) 2.5 µl, 20
 pmole forward and turn around groundwork
 each 2.0 µl, dNTP blend (2.5mM) 3.0 µl, 0.5
 µl of Taq DNA polymerase (2.5 U),
 nuclease free water and 50 ng of DNA
 format. DNA tests were opened up on DNA
 thermalcycler (T 100, BioRad, USA). The
 PCR condition was as per the following:
 beginning denaturation for 3 min at 94°C,
 30 cycles each comprising of denaturation
 for 1 min at 94°C, preliminary strengthening
 for 1 min at 54 °C and augmentation at 72°C
 for 5 min and, a last prolongation of 5 min at
 72°C. An aliquot of 3-4 µl of PCR response
 item was electrophoresed on a 1% agarose
 gel containing ethidium bromide (0.5µg ml⁻¹
 in H₂O) and the DNA groups were
 envisioned under the UV light in a gel
 documentation framework (BIORAD,
 India).

**Sequencing and phylogenetic
 investigation**

Amplicon of size 1.5 KB was filtered
 utilizing a DNA cleaning pack (Quiagen,
 USA) and sequencing was performed at
 Xcelris Genomics Labs Ltd. (Xcelris, India).
 The sequenced nucleotides were looked at
 against the GenBank database utilizing the
 NCBI BLAST calculated and stored in the
 NCBI information base
<http://www.ncbi.nlm.nih.gov/BLAST>. An
 ordered association of confines was
 relegated utilizing RDP
 database(http://rdp.cme.msu.edu/seqmatch/seqmatch_intro.jsp) at 98% limit of 16S
 rRNA quality grouping. A Phylogenetic
 examination was finished by utilizing
 Software MEGA 6.0 (Tamura et al., 2013)
 and adjusted utilizing CLUSTAL-X. The
 pairwise transformative separation was
 developed by the Neighbor Joining
 technique with the bootstrap of 500 imitates
 to group the related .

**Plant development advancing test ACC
 deaminase examine**

Bacterial disconnect was developed in Tryptic soya soup (Himedia, India) to late log stage at 200 rpm for 24 h at 30 °C. The gathered cells were reaped by centrifugation, washed with 0.1M Tris HCl (pH 7.6) and hatched overnight in 7.5 ml insignificant medium containing 3 mM ACC as sole nitrogen source. The bacterial cells were come back to shaking water shower for acceptance of chemical at 200 rpm for 24 h at 30 °C. At that point cells were gathered by centrifugation, washed subsequently with 0.1M Tris-HCl (pH 7.6) and resuspended in 0.1 M Tris-HCl (pH 8.5). At long last thirty μ l toluene was added to cell suspension and homogenized for 30 s. Now 100 μ l of toluenized cells were put aside for protein examine and put away at 4 °C.

Plant growth promoting test

Disengaged bacterium was screened for different plant development advancing (PGP) highlights in charge of plant development advancement. On measurement, ACC deaminase action of

confine was observed to be 157.00 ± 2.7 nmol μ g⁻¹Pr.h⁻¹. In light of the unmistakable zone saw around the streaked settlement of bacterium on tricalcium phosphate plate, it was recognized as positive for the trial of phosphate (P) solubilization.

Antagonistic test

Biocontrol potential of strain SNP-18 was evaluated by testing antagonistic activity against various fungal and bacterial pathogens. This isolate was observed to be inhibitory for the growth of *Fusarium oxysporum* (13.1 ± 0.21) and *Penicillium citrium* (14.8 ± 0.42) whereas ineffective against *Aspergillus flavus*, *Fusarium graminearum* and *Fusarium moniliforme*.

Stress tolerance test

Isolated bacterial strain SNP-18 was tested for its ability to tolerate abnormal growth conditions by incubating it to a wide range of salt, pH and temperature stress condition.

Following experimental observation, isolate was found to tolerate salt concentration in range of 4% to 6%. The pH tolerance of isolate was in range of pH 5.0 to 11.0. In addition, isolate can grow up to higher temperature of 50 °C.

Discussion

Plant rhizosphere is a favored specialty for soil microorganisms because of rich supplement accessibility. Different root exudates discharged by plants draw in various genera of microbial species. Assorted microorganism having a place with genera like Rhizobium, Msorhizobium, Bradyrhizobium, Azotobacter, Azospirillum and Enterobacter improved the plant development thus of their air nitrogen obsession capacity. In any case, regardless of nitrogen obsession, development advancement of plants might be ascribed to different instruments, for example, generation of plant development directing phytohormones, IAA creation and ACC deaminase action (Arshad and

Frankenberger, 1993; Glick, 1995). The creation of indole acidic corrosive shifts with bacterial secludes and centralization of tryptophan, albeit characteristic capacity of microorganism to deliver IAA in the rhizosphere relies on upon the accessibility of forerunners and take-up of microbial IAA by plant (Arshad and Frankenberger, 1993). Another vital attribute of plant development advancing rhizospheric microorganism that may by implication impact the plant development, is the creation of siderophores. However in the present review disconnect was discovered negative for siderophore creation. They tie to the accessible type of iron Fe^{3+} in the rhizosphere and limit its accessibility to phytopathogens and subsequently assume a key part in upkeep of plant wellbeing. The antifungal movement of the test disengages demonstrated a cozy association with the generation of HCN or synergistic cooperation with different metabolites.

Conclusion

Microorganism with plant development advancing characteristics will be useful to build trim profitability under anxiety condition. Our discoveries fundamentally concentrate on detachment of viable anxiety tolerant microscopic organisms that have proficient ACCD movement to diminish stretch initiated ethylene level in plants, subsequently could be utilized as a compelling instrument to build the development and profitability for plants development. In a word, watched brings about this review show that microbes with various PGP characteristics will be more gainful than having single qualities. Consequently utilization of such diverse development advancing microbes could be utilized as biofertilizers for plants developing under anxiety situations.

REFERENCES:

- Arshad M and Frankenberger Jr WT (1993). Microbial production of plant growth regulators. In: Blaine, F. Metting, Jr. (Eds.), Soil Microbial Ecology. Marcel and Dekker, Inc., New York, pp. 307–347.
- Cappuccino JC and Sherman N (1992). In: Microbiology: A Laboratory Manual, third ed. Benjamin/cummings Pub.Co., New York, pp. 125–179.
- Cattelan AJ, Hartel PG, Fuhrmann JJ (1999). Screening of plant growth_promoting rhizobacteria to promote early soybean growth. Soil Sci. Soc. Am. J. 63, 1670–1680.
- Dobereiner J (1995). Isolation and identification of aerobic nitrogen fixing bacteria from soil and plants. In: Alef, K. and P. Nanniperi (eds.), Methods Appl. Soil Microbiol. Biochem. pp: 134– 141. Academic press, London.
- Fuhrmann JJ, Wollum II AG (1989). Nodulation competition among Bradyrhizobium japonicum strains

- influenced by rhizosphere bacteria and iron availability. *Biol. Fertil. Soil.* 7, 108–112.
- Gaur AC (1990). Physiological functions of phosphate solubilizing micro-organisms. In: Gaur, A.C. (Ed.), *Phosphate Solubilizing Micro-organisms as Biofertilizers*. Omega Scientific Publishers, New Delhi, pp. 16–72.
- Glick BR (1995). The enhancement of plant growth by free living bacteria. *Can. J. Microbiol.* 41, 109–114. Glick BR (2007). Promotion of plant growth by bacterial ACC deaminase. *Crit. Rev. Plant Sci.* 26, 227–242.
- Gordon SA and Weber RP (1951). Colorimetric estimation of indole acetic acid. *Plant Physiol.* 26, 192–195.
- antarum 118, 10-15.
- Holbrook A, Edge W, Bailey F (1961). Spectrophotometric method for determination of gibberellic acid. *Adv. Chem. Ser.* 28, 159–167.
- Nagrajkumar M, Bhaaskaran R, Velazhahan R (2004). Involvement of secondary metabolites and extracellular lytic enzymes produced by *Pseudomonas fluorescens* in inhibition of *Rhizoctonia solani*, the rice sheath of blight pathogen. *Microbiol. Res.* 159, 73–81.
- Penrose DM and Glick BR (2003). Methods for isolating and characterizing ACC deaminase-containing plant growth-promoting rhizobacteria. *Physiologia Pl*